MISC-2012-14

Is 125 GeV techni-dilaton found at LHC?

Shinya Matsuzaki[1, *] and Koichi Yamawaki[2, †]

[1] *Maskawa Institute for Science and Culture, Kyoto Sangyo University,*
Motoyama, Kamigamo, Kita-Ku, Kyoto 603-8555, Japan.
[2] *Kobayashi-Maskawa Institute for the Origin of Particles and the Universe (KMI)*
Nagoya University, Nagoya 464-8602, Japan.

(Dated: July 26, 2012)

A new particle at around 125 GeV has been observed at the LHC, which we show could be identified with the techni-dilaton (TD) predicted in the walking technicolor and thus should be an evidence of walking technicolor. The TD is a pseudo Nambu-Goldstone boson for the approximate scale symmetry spontaneously broken by techni-fermion condensation, with its lightness being ensured by the approximate scale invariance of the walking technicolor. We test the goodness-of-fit of the TD signatures using the presently available LHC data set, and show that the 125 GeV TD is actually favored by the current data to explain the reported signal strengths in the global fit as well as in each channel including the coupling properties, most notably the somewhat large diphoton event rate. This is in sharp contrast to other dilaton/radion scenarios which tend to be disfavored by the data, mainly due to the suppressed diphoton event rate.

arXiv:1207.5911v1 [hep-ph] 25 Jul 2012

I. INTRODUCTION

On July 4, the ATLAS and CMS groups [1] have reported observation of a new boson at around 125 GeV in search for the decay channels such as $\gamma\gamma$, WW^* and ZZ^*. Though the statistical uncertainty is still large, the current data on the diphoton event rate [2, 3] exhibit the signal strength about two times larger than that expected from the standard model (SM) Higgs boson, which may imply the observation of a new boson beyond the SM.

One such a possibility is the techni-dilaton (TD), a composite scalar boson, predicted in the walking technicolor (WTC) [4, 5] which is characterized by an approximately scale-invariant (conformal) gauge dynamics and a large anomalous dimension $\gamma_m = 1$ (The WTC was subsequently studied without notion of anomalous dimension and scale invariance/TD. [6]). The TD is a pseudo Nambu-Goldstone boson for the spontaneous breaking of the approximate scale symmetry triggered by techni-fermion condensation and hence its lightness, say 125 GeV, is protected by the approximate scale symmetry inherent to the WTC. Thus the discovery of TD would imply the discovery of the WTC.

In Refs. [7–9] we studied the LHC signatures of the TD. Particularly in Ref. [9] we have shown that the 125 GeV TD can be consistent with the currently reported diphoton signal as well as other signals such as WW^* and ZZ^*, etc..

In this article, we extend the previously reported analysis [9] on the 125 GeV TD by testing the goodness-of-fit of the TD signatures, based on the presently available LHC data set [2, 3]. We show that the 125 GeV TD is actually favored (slightly better than the SM Higgs) by testing the goodness-of-fit of the current data: The large diphoton event rate is achieved due to the presence of extra techni-fermion loop contributions, a salient feature of the techni-dilaton noted in the previous papers [7–9]. It is also shown that, lacking such extra fermion contributions, a similar dilaton model [10] and a typical radion model [11] analyzed recently for LHC [12, 13] tend to be disfavored by the data, mainly due to the suppressed diphoton rate.

II. TD COUPLINGS

The TD couplings to techni-fermions and the SM particles have been discussed in detail in Ref. [9], which are completely fixed by Ward-Takahashi identities for the dilatation current coupled to the TD. Those couplings are collected in an effective nonlinear Lagrangian invariant under the scale symmetry [9]:

$$
\begin{aligned}
\mathcal{L} \;=\; & \frac{F_\pi^2}{4}\chi^2 \mathrm{Tr}[\mathcal{D}_\mu U^\dagger \mathcal{D}^\mu U] + \frac{F_\phi^2}{2}(\partial_\mu \chi)^2 \\
& - m_f \left(\left(\frac{\chi}{S}\right)^{2-\gamma_m} \cdot \chi \right) \bar{f} f \\
& + \log\left(\frac{\chi}{S}\right) \left\{ \frac{\beta_F(g_s)}{2g_s} G_{\mu\nu}^2 + \frac{\beta_F(e)}{2e} F_{\mu\nu}^2 \right\} + \cdots , \qquad (1)
\end{aligned}
$$

where $\chi = e^{\phi/F_\phi}$ with the TD field ϕ and decay constant F_ϕ, and $U = e^{2i\pi/F_\pi}$ with the techni-pion fields π and decay constant F_π related to the electroweak (EW) scale $v_{\rm EW} \simeq 246$ GeV by $F_\pi = v_{\rm EW}/\sqrt{N_D}$, in which N_D denotes the number of EW doublets formed by the techni-fermions. The SM gauge covariant $\mathcal{D}_\mu U$ term includes the weak gauge boson masses in the usual manner. A spurion field S has been introduced so as to compensate the scale invariance explicitly broken by the techni-fermion condensation itself, where S is set to 1 after all calculations as in the case of other spurion methods.

---

[*] `synya@cc.kyoto-su.ac.jp`

[†] `yamawaki@kmi.nagoya-u.ac.jp`

The TD Yukawa coupling to the SM f-fermion arises from the second line of Eq.(1) as [5],

$$-\frac{(3-\gamma_m)m_f}{F_\phi}\,\phi\bar{f}f\,, \tag{2}$$

along with scale dimension of techni-fermion bilinear operator $(3-\gamma_m)$, where the anomalous dimension is $\gamma_m \simeq 1$ in WTC, which is crucial to obtain the realistic mass of the SM fermions of the first and the second generations without suffering from the FCNC (flavor-changing neutral current) problems [14]. However it was known for long time that it is not enough for the mass of the third-generation SM f-fermions like t, b, τ: A simplest resolution would be the strong extended-technicolor (ETC) TC model [15] having much larger anomalous dimension $1 < \gamma_m < 2$ due to the strong effective four-fermion coupling from the ETC dynamics in addition to the walking gauge coupling. As was prescribed in *Note added* of Ref. [9], here we take $\gamma_m \simeq 2$, i.e., $(3-\gamma_m) \simeq 1$, as in the strong ETC model [15] for the third-generation SM f-fermions like t, b, τ which are relevant to the current LHC data.

From the Lagrangian Eq.(1) one thus easily sees that the TD couplings to W and Z bosons and fermions are related to those of the SM Higgs by a simple scaling:

$$\begin{aligned} \frac{g_{\phi WW/ZZ}}{g_{h_{\rm SM}WW/ZZ}} &= \frac{v_{\rm EW}}{F_\phi}\,, \\ \frac{g_{\phi ff}}{g_{h_{\rm SM}ff}} &= \frac{v_{\rm EW}}{F_\phi}\,, \qquad \text{for} \quad f = t, b, \tau\,. \end{aligned} \tag{3}$$

In addition to the above scaling, the couplings to gluon and photon ($G_{\mu\nu}^2$ and $F_{\mu\nu}^2$ terms in Eq.(1)) involve the beta functions, $\beta_F(g_s)$ and $\beta_F(e)$, induced from F-techni-fermion loops [7]. This is the most crucial point [9] that distinguishes the TD from other similar models of dilaton/radion [10, 11]. Here we shall employ the one-family model (1FM) with $N_D = 4$ as a WTC setting (see the later discussions) for the $SU(N_{\rm TC})$ gauge group. The beta functions $\beta_F(g_s)$ and $\beta_F(e)$ are then explicitly evaluated to be

$$\begin{aligned} \beta_F(g_s) &= \frac{g_s^3}{(4\pi)^2}\frac{4}{3}N_{\rm TC} \\ \beta_F(e) &= \frac{e^3}{(4\pi)^2}\frac{16}{9}N_{\rm TC}\,. \end{aligned} \tag{4}$$

We thus find the scaling from the SM Higgs for the couplings to gg and $\gamma\gamma$, which can approximately be expressed at around 125 GeV as

$$\begin{aligned} \frac{g_{\phi gg}}{g_{h_{\rm SM}gg}} &\simeq \frac{v_{\rm EW}}{F_\phi}\cdot\left((3-\gamma_m)+2N_{\rm TC}\right) \\ \frac{g_{\phi\gamma\gamma}}{g_{h_{\rm SM}\gamma\gamma}} &\simeq \frac{v_{\rm EW}}{F_\phi}\cdot\left(\frac{63-16(3-\gamma_m)}{47}\frac{32}{47}N_{\rm TC}\right)\,, \end{aligned} \tag{5}$$

where in estimating the SM contributions we have incorporated only the top and W boson loop contributions. Note that since we used $\gamma_m = 2$ for the top quark Yukawa coupling as noted before (as well as *Note added* of Ref. [9]), Eq.(5) is slightly different from Eq. (51) of [9] (which used $\gamma_m = 1$) for the numerical analysis in the present work. The full expressions for these couplings are presented in Appendix A of Ref. [9].

As seen from Eqs.(3) and (5), once the ratio $v_{\rm EW}/F_\phi$ is fixed, the TD LHC phenomenological study can be made just by quoting the SM Higgs coupling properties. The TD decay constant F_ϕ can actually be related to the TD mass M_ϕ through the partially conserved dilatation current (PCDC) – which is analogous to the PCAC (partially conserved axialvector current) relation for the QCD pion – involving the vacuum energy density $\mathcal{E}_{\rm vac}$:

$$F_\phi^2 M_\phi^2 = -16\mathcal{E}_{\rm vac}\,. \tag{6}$$

The vacuum energy density $\mathcal{E}_{\rm vac}$ is saturated by the techni-gluon condensation induced from the techni-fermion condensation, so is generically expressed as

$$\mathcal{E}_{\rm vac} = -\kappa_V\left(\frac{N_{\rm TC}N_{\rm TF}}{8\pi^2}\right)m_F^4\,, \tag{7}$$

where m_F denotes the dynamical techni-fermion mass and $N_{\rm TF} = 2N_{\rm D} + N_{\rm EW-singlet}$ including the number of dummy techni-fermions, $N_{\rm EW-singlet}$, which are singlet under the SM gauges. The overall coefficient κ_V is determined once a straightforward nonperturbative calculation is made. The dynamical techni-fermion mass m_F can, on the other hand, be related to the techni-pion decay constant F_π:

$$F_\pi^2 = \kappa_F^2\frac{N_{\rm TC}}{4\pi^2}m_F^2\,, \tag{8}$$

with the overall coefficient κ_F and the property of $N_{\rm TC}$ scaling taken into account.

The values of κ_V and κ_F may be quoted from the latest result [16] on a ladder Schwinger-Dyson analysis for a modern version of WTC [17–19]:

$$\kappa_V \simeq 0.7\,, \qquad \kappa_F \simeq 1.4\,, \tag{9}$$

where κ_F has been estimated based on the Pagels-Stokar formula [20]. In that case $N_{\rm TF}$ is fixed by the criticality condition for the walking regime as [18]

$$\frac{N_{\rm TF}}{4N_{\rm TC}} \simeq 1\,. \tag{10}$$

The estimated values in Eqs.(9) and (10) are based on ladder approximation which are subject to certain uncertainties up to 30% observed for the critical coupling and hadron spectrum in QCD [21]. We may include this 30% uncertainty in estimation of each independent factor κ_V, κ_F^2 and the criticality condition $N_{\rm TF}/(4N_{\rm TC})$. Putting these all together, we thus estimate $v_{\rm EW}/F_\phi$ as

$$\frac{v_{\rm EW}}{F_\pi} \simeq (0.1-0.3)\times\left(\frac{N_D}{4}\right)\left(\frac{M_\phi}{125\,{\rm GeV}}\right)\,, \tag{11}$$

and $m_F \simeq (320 - 420)\mathrm{GeV}\sqrt{3/N_{\rm TC}}$ for the 1FM with $F_\pi \simeq 123$ GeV.

From Eqs.(3) and (11) we see that *the TD couplings to* WW, ZZ *and* $f\bar{f}$ *are substantially smaller than those of the SM Higgs. On the other hand, the TD couplings to* gg *and* $\gamma\gamma$ *in Eq.(5) have extra factors* $(1 + 2N_{\rm TC})$ *and* $(1 - 32N_{\rm TC}/47)$ *coming from techni-fermions as well as the* W *and top quarks carrying the QCD color and electromagnetic charges.* The gluon fusion (GF) production at the LHC thus becomes larger than the SM Higgs case due to this extra factor, while other productions such as vector boson fusion (VBF) and vector boson associate (VBA) productions are significantly suppressed.

III. TD SIGNAL AT LHC

As done in Refs. [8, 9], we can calculate the TD production cross sections and decay widths including loop contributions from the SM particles, by quoting the corresponding formulas for the SM Higgs [22]. (The explicit expressions of the formulas for TD are given in Ref. [9].) Here we focus on the GF and VBF productions for decay channels to WW^*, ZZ^*, $\tau^+\tau^-$ and $\gamma\gamma$, and VBA production for $b\bar{b}$ channel, to which the ATLAS and CMS experiments have so far reported the significant data in search for the SM Higgs [2, 3].

The signal strength $\mu \equiv \sigma/\sigma_{\rm SM}$ for $b\bar{b}$ channel at $\sqrt{s} = 7, 8$ TeV is thus evaluated as

$$\begin{aligned} \mu_{b\bar{b}} &= \frac{\sigma^{\phi}_{\rm VBA}(s)}{\sigma^{h_{\rm SM}}_{\rm VBA}(s)} \frac{BR(\phi \to b\bar{b})}{BR(h_{\rm SM} \to b\bar{b})} \\ &= \frac{\sigma_{W\phi}(s) + \sigma_{Z\phi}(s)}{\sigma_{Wh_{\rm SM}}(s) + \sigma_{Zh_{\rm SM}}(s)} \frac{BR(\phi \to b\bar{b})}{BR(h_{\rm SM} \to b\bar{b})} \,. \end{aligned} \tag{12}$$

For $X = WW^*$, ZZ^* and $\tau^+\tau^-$ channels we take the signal strengths to be inclusive:

$$\mu_X = \frac{\sigma^{\phi}_{\rm GF}(s) + \sigma^{\phi}_{\rm VBF}(s)}{\sigma^{h_{\rm SM}}_{\rm GF}(s) + \sigma^{h_{\rm SM}}_{\rm VBF}(s)} \frac{BR(\phi \to X)}{BR(h_{\rm SM} \to X)} \,. \tag{13}$$

On the other hand, $\gamma\gamma + 0j$ and $\gamma\gamma + 2j$ channels are treated to be exclusive by distinguishing the TD production processes:

$$\begin{aligned} \mu_{\gamma\gamma 0j} &= \frac{\sigma^{\phi}_{\rm GF}(s)}{\sigma^{h_{\rm SM}}_{\rm GF}(s)} \frac{BR(\phi \to X)}{BR(h_{\rm SM} \to X)} \,, \\ \mu_{\gamma\gamma 2j} &= \frac{\xi_{\rm GF} \cdot \sigma^{\phi}_{\rm GF}(s) + \xi_{\rm VBF} \cdot \sigma^{\phi}_{\rm VBF}(s)}{\xi_{\rm GF} \cdot \sigma^{h_{\rm SM}}_{\rm GF}(s) + \xi_{\rm VBF} \cdot \sigma^{h_{\rm SM}}_{\rm VBF}(s)} \\ &\quad \times \frac{BR(\phi \to \gamma\gamma)}{BR(h_{\rm SM} \to \gamma\gamma)} \,, \end{aligned} \tag{14}$$

where the corresponding acceptances multiplied by dijet tag efficiencies $\xi_{\rm GF}$ and $\xi_{\rm VBF}$ are read off from Refs. [2, 3]. We then combine the 7 TeV and 8 TeV signal strengths with the luminosities accumulated for each event category. It turns out that $\mu_{\gamma\gamma 0j}$ can be enhanced by the factors both from the gg and $\gamma\gamma$ couplings (See Eq.(5)), which can compensate the smallness of $(v_{\rm EW}/F_\phi)$ in Eq.(11) for a moderately large $N_{\rm TC}$ [9].

This feature is in sharp contrast to other similar dilaton models such as EW pseudo-dilaton [10] and Randall-Sundrum (RS) radion [11], where extra contributions beyond the SM such as techni-fermions are absent so that their diphoton rates are not enhanced, to be disfavored by the current diphoton data as will be shown more explicitly below.

We shall test the goodness-of-fit of the TD, based on the χ^2 function:

$$\chi^2 = \sum_{i \in \text{events}} \left(\frac{\mu_i - \mu_i^{\rm exp}}{\sigma_i} \right)^2 \,, \tag{15}$$

where $\mu_i^{\rm exp}$ denote the best-fit strengths for each channel reported in Refs [2, 3] and σ_i the corresponding one sigma errors. Taking $(v_{\rm EW}/F_\phi)$ as a free parameter so as to satisfy the theoretically expected range in Eq.(11), in Fig. 1 we plot the χ^2 function for the 125 GeV TD in the 1FM with $N_{\rm TC} = 4, 5$. The best-fit values are as follows:

$N_{\rm TC}$	$(v_{\rm EW}/F_\phi)_{\rm best}$	$\chi^2_{\rm min}$/d.o.f
4	0.22	$12/13 \simeq 0.9$
5	0.17	$10/13 \simeq 0.7$

, (16)

which are compared with the SM Higgs case, $\chi^2_{\rm min}|_{\rm SM} \simeq 14/14 = 1.0$, implying that the TD is more favorable than the SM Higgs ($\mu_i = 1$). This nice goodness of fit is due to the significant enhancement in the diphoton channel coming from the sector beyond the SM (techni-fermions), as was also noted in Ref. [23]: The techni-fermion loop contributions as in Eq.(5) become large enough to compensate the smallness of the overall $(v_{\rm EW}/F_\phi)$ in Eq.(11).

In Fig. 1 also has been shown a comparison with other similar models, EW pseudo-dilaton [10] and RS radion [11]. These dilaton/radion scenarios are actually disfavored mainly due to the absence of enhancement of diphoton rate, in sharp contrast to the TD [24]. This result is in accord with other similar analyses in Ref. [25] for the EW pseudo-dilaton and in Ref. [26] for the RS radion performed in light of the 125 GeV LHC events. although we have not included the Tevatron results on the $\bar{b}b$ channel [27].

Finally, in Fig. 2 we explicitly compare the best-fit signal strengths of TD with those estimated by the ATLAS and CMS analyses for each channel. Note first that the most precise measurement has currently been done in the diphoton channel with 0 jet ($\gamma\gamma 0j$) which exhibits about 2 times larger signal strengths than the SM Higgs prediction ($\mu_{\gamma\gamma 0j} = 1.9 \pm 0.5$ for ATLAS 7TeV + 8TeV and $\mu_{\gamma\gamma 0j} = 1.7 \pm 0.5$ for CMS 7TeV + 8TeV [2, 3]). The χ^2 fit is therefore fairly sensitive to the $\gamma\gamma 0j$ category, and hence currently the TD can be more favorable than the SM Higgs due to the enhancement of the diphoton rate which happens when $N_{\rm TC} \geq 4$. On the other hand, the TD signal strength in the diphoton plus dijet channel

($\gamma\gamma 2j$) tends to be smaller than the SM Higgs prediction, simply because of the suppression of the overall TD coupling compared to the SM Higgs ($v_{\rm EW}/F_\phi$) in Eq.(11). Similar suppressions are also seen in other exclusive channels like $2l2\nu + 2j$ and $\tau^+\tau^- + 2j$ as well as $b\bar{b}$ originated from the VBA and VBF productions. Thus more precise measurements in such other exclusive events would draw a more definite conclusion that the TD is favored or not.



FIG. 1: The plot of χ^2 as a function of $v_{\rm EW}/F_\phi$ in the case of the 1FM with $N_{\rm TC} = 4$ (black dashed curve) and $N_{\rm TC} = 5$ (black solid curve). Comparison with the EW pseudo-dilaton [10] (blue dotted curve) and RS radion [13] (green dotted curve) is also shown, along with the SM Higgs case (red dotted line). Here $\chi^2_{\rm min}/{\rm d.o.f} = 12/13(10/13) \simeq 0.9(0.7)$ for the TD with $N_{\rm TC} = 4(5)$, $\chi^2_{\rm min}/{\rm d.o.f} = 37/13 \simeq 2.8$ for the EW pseudo-dilaton and $\chi^2_{\rm min}/{\rm d.o.f} = 30/13 \simeq 2.3$ for the RS radion.



FIG. 2: The best-fit signal strengths of the 125 GeV TD, for the decay channels categorized as $WW^*(2l2\nu)$, $ZZ^*(4l)$, $\tau^+\tau^-$, $\gamma\gamma 0j$ and $\gamma\gamma 2j$ [2, 3].

IV. SUMMARY

In summary, we have tested the goodness-of-fit of the 125 GeV TD signatures, using the presently available LHC data set by focusing on the decay channels to $b\bar{b}$, WW^*, ZZ^*, $\tau^+\tau^-$ and $\gamma\gamma$. We found that the 125 GeV TD is actually favored by the current data to explain the reported signal strengths including the coupling properties involving somewhat large diphoton event rate. It was also shown that other dilaton and radion scenarios tend to be disfavored by the data, mainly due to the suppressed diphoton event rate.

The issue to be studied in future is a reliable nonperturbative estimate of the TD mass: There has been only a crude estimate based on the ladder approximation (see references cited in e.g. Ref.[9]). The holographic method [28] may give some hint and the lattice simulations eventually give us the definite answer.

Acknowledgments

This work was supported by the JSPS Grant-in-Aid for Scientific Research (S) #22224003 and (C) #23540300 (K.Y.).

[1] J. Incandela, CMS talk at "*Latest update in the search for the Higgs boson at CERN*", July 4, 2012; F. Gianotti, ATLAS talk at "*Latest update in the search for the Higgs boson at CERN*", July 4, 2012.

[2] The ATLAS Collaboration, ATLAS-CONF-2012-014; ATLAS-CONF-2012-019; ATLAS-CONF-2012-091; ATLAS-CONF-2012-092; ATLAS-CONF-2012-093; ATLAS-CONF-2012-098.

[3] The CMS Collaboration, CMS-PAS-HIG-12-015; CMS-PAS-HIG-12-016; CMS-PAS-HIG-12-017; CMS-PAS-HIG-12-018; CMS-PAS-HIG-12-019; CMS-PAS-HIG-12-020.

[4] K. Yamawaki, M. Bando and K. Matumoto, Phys. Rev. Lett. **56**, 1335 (1986); M. Bando, T. Morozumi, H. So and K. Yamawaki, Phys. Rev. Lett. **59**, 389 (1987).

[5] M. Bando, K. Matumoto and K. Yamawaki, Phys. Lett. B **178**, 308 (1986).

[6] T. Akiba and T. Yanagida, Phys. Lett. B **169**, 432 (1986); T. W. Appelquist, D. Karabali and L. C. R. Wijewardhana, Phys. Rev. Lett. **57**, 957 (1986). A similar analysis was done earlier on the purely numerical basis by B. Holdom, Phys. Lett. B **150**, 301 (1985).

[7] S. Matsuzaki and K. Yamawaki, Prog. Theor. Phys. **127**, 209 (2012).

[8] S. Matsuzaki and K. Yamawaki, Phys. Rev. D **85**, 095020 (2012).

[9] S. Matsuzaki and K. Yamawaki, arXiv:1206.6703 [hep-ph].

[10] W. D. Goldberger, B. Grinstein and W. Skiba, Phys. Rev. Lett. **100**, 111802 (2008); J. Fan, W. D. Goldberger, A. Ross and W. Skiba, Phys. Rev. D **79**, 035017 (2009); L. Vecchi, Phys. Rev. D **82**, 076009 (2010);

[11] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999); **83**, 4690 (1999); W. D. Goldberger and M. B. Wise, Phys. Rev. Lett. **83**, 4922 (1999).

[12] B. Coleppa, T. Gregoire and H. E. Logan, Phys. Rev. D **85**, 055001 (2012); V. Barger, M. Ishida and W. -Y. Keung, Phys. Rev. D **85**, 015024 (2012); Phys. Rev. Lett. **108**, 101802 (2012); B. A. Campbell, J. Ellis and K. A. Olive, JHEP **1203**, 026 (2012).

[13] V. Barger and M. Ishida, Phys. Lett. B **709**, 185 (2012).

[14] Other issues such as the S and T parameters were discussed in e.g., Ref. [9].

[15] V. A. Miransky, K. Yamawaki, Mod. Phys. Lett. **A4**, 129-135 (1989); K. Matumoto, Prog. Theor. Phys. **81**, 277-279 (1989); T. Appelquist, M. Einhorn, T. Takeuchi, L. C. R. Wijewardhana, Phys. Lett. **B220**, 223 (1989).

[16] M. Hashimoto and K. Yamawaki, Phys. Rev. D **83**, 016008 (2011).

[17] K. D. Lane and M. V. Ramana, Phys. Rev. D **44**, 2678 (1991).

[18] T. Appelquist, J. Terning and L. C. Wijewardhana, Phys. Rev. Lett. **77**, 1214 (1996); T. Appelquist, A. Ratnaweera, J. Terning and L. C. Wijewardhana, Phys. Rev. D **58**, 105017 (1998).

[19] V. A. Miransky and K. Yamawaki, Phys. Rev. D **55**, 5051 (1997); Errata, **56**, 3768 (1997).

[20] H. Pagels and S. Stokar, Phys. Rev. D **20**, 2947 (1979).

[21] T. Appelquist, K. D. Lane and U. Mahanta, Phys. Rev. Lett. **61**, 1553 (1988); K. -I. Aoki, T. Kugo and M. G. Mitchard, Phys. Lett. B **266**, 467 (1991); M. Harada, M. Kurachi and K. Yamawaki, Phys. Rev. D **70**, 033009 (2004).

[22] M. Spira, Fortsch. Phys. **46**, 203 (1998) [arXiv:hep-ph/9705337]; S. Dittmaier *et al.* [LHC Higgs Cross Section Working Group Collaboration], arXiv:1101.0593 [hep-ph]; LHC Higgs Cross Section Working Group *et al.*, arXiv:1201.3084 [hep-ph]; *https://twiki.cern.ch/twiki/bin/view/LHCPhysics /CERNYellowReportPageAt8TeV.*

[23] J. Ellis and T. You, arXiv:1207.1693 [hep-ph].

[24] A variant of the RS radion such as the one discussed in Ref. [29] could actually be as favorable as the TD because of an accidental enhancement of the diphoton event rate induced solely from a choice of the SM sector contributions.

[25] D. Carmi, A. Falkowski, E. Kuflik, T. Volansky and J. Zupan, arXiv:1207.1718 [hep-ph]; I. Low, J. Lykken and G. Shaughnessy, arXiv:1207.1093 [hep-ph].

[26] P. P. Giardino, K. Kannike, M. Raidal and A. Strumia, arXiv:1207.1347 [hep-ph].

[27] Even including the Tevatron results for $b\bar{b}$, WW^* and $\gamma\gamma$ channels, the goodness-of-fit of TD does not substantially change so much ($\chi^2_{\rm min}$/d.o.f $= 19/16 \simeq 1.2$ compared with the SM Higgs case, $\chi^2_{\rm min}$/d.o.f $= 18/17 \simeq 1.1$).

[28] K. Haba, S. Matsuzaki and K. Yamawaki, Phys. Rev. D **82**, 055007 (2010); S. Matsuzaki and K. Yamawaki, in preparation; D. Kutasov, J. Lin and A. Parnachev, Nucl. Phys. B **863**, 361 (2012); R. Lawrance and M. Piai, arXiv:1207.0427 [hep-ph].

[29] K. Cheung and T. -C. Yuan, Phys. Rev. Lett. **108**, 141602 (2012) [arXiv:1112.4146 [hep-ph]].